

Retrospective comparison of external beam radiotherapy and radical prostatectomy in high-risk, clinically localized prostate cancer: Why EBRT + ADT remains the gold standard



C. Gonzalez-San Segundo¹, F. Herranz Amo², G. Ogaya Pinies², P. Cuesta Álvaro³, F. Calvo Manuel¹, J. Santos-Miranda¹

¹ Hospital General Universitario Gregorio Marañón, Oncología Radioterápica, Spain

² Hospital General Universitario Gregorio Marañón, Urología, Spain

³ Universidad Complutense de Madrid, Informatics Department for Research Support, Spain

Purpose. To retrospectively compare results of radical prostatectomy (RP) without adjuvant treatment versus radical external beam radiotherapy (EBRT) plus androgen deprivation therapy (ADT) for high-risk prostate cancer (PCa) patients.

Methods and materials. Between 1995 and 2008, 305 patients with high-risk PCa were treated at the Urology and Oncology departments. 140 patients underwent open RP (74% with lymph node dissection) and 165 underwent definitive EBRT (86% with ADT). 95% received total dose ≥ 74 Gy. The 2 groups were homogeneous in terms of cT stage, Gleason score and initial PSA level. Biochemical relapse was defined as a PSA level > 0.4 ng/ml for RP patients and nadir + 2 for EBRT. The endpoint selected for analysis was to compare biochemical recurrence-free survival (bRFS) among groups.

Results. The median follow-up was 74 months [8–184]. The 7-year bRFS rates for RP and EBRT patients were $41.5 \pm 4\%$ and $80 \pm 4\%$, respectively ($p = 0.00$). The disease-specific survival at 7 years was $91 \pm 2\%$ in the surgical group and $95 \pm 2\%$ in the EBRT cohort ($p = ns$). Seventy-two of 81 patients with BF after surgery (24 of them had persistent high PSA), were treated with EBRT salvage \pm ADT (51% of the surgical patients). In 22 of 29 patients with BF after EBRT (13% of the irradiated group), salvage treatment with ADT or chemotherapy was administered. Only 1 patient was treated with local treatment. At the time of analysis, a total of 30 patients had cancer progression, 20 in the surgery group and 10 in EBRT + ADT group (clinical progression-free survival at 7 years, 85% and 94%, respectively).

Conclusions. Improved outcome with EBRT is observed compared to RP in patients with high-risk PCa regarding bRFS. However, specific cancer mortality was similar between PR and EBRT groups, which emphasizes the efficacy of salvage therapy after surgery.

<http://dx.doi.org/10.1016/j.rpor.2013.03.823>

Testosterone kinetics after androgen-deprivation therapy in intermediate and high risk prostate cancer: Results from a randomized trial (DART01/05)



C. González-san Segundo¹, A. Zapatero², A. Guerrero³, X. Maldonado⁴, A. Álvarez¹, M. Cabeza⁵, V. Macías⁶, F. Casas⁷, A. Pedro-olivé⁸, A. Boladeras⁹, C. Martín de Vidales², M. Vázquez¹⁰, F. Calvo¹

¹ Hospital General Universitario Gregorio Marañón, Oncología Radioterápica, Spain

² Hospital Universitario de la Princesa, Oncología Radioterápica, Spain

³ Complejo Asistencial Son Dureta, Oncología Radioterápica, Spain

⁴ Hospital General Universitari Vall D'hebron, Oncología Radioterápica, Spain

⁵ Hospital 12 de Octubre, Oncología Radioterápica, Spain

⁶ Hospital Universitario de Salamanca. H. Virgen de la Vega, Oncología Radioterápica, Spain

⁷ Hospital Clínic I Provincial de Barcelona, Oncología Radioterápica, Spain

⁸ Clínica Plató, Fundació Privada, Oncología Radioterápica, Spain

⁹ Institut Català D'oncologia, Oncología Radioterápica, Spain

¹⁰ Hospital do Meixoeiro, Oncología Radioterápica, Spain

Background. Limited information exists on the absolute relationship between testosterone (T) values and clinical outcomes in prostate cancer (PCa).

Purpose. To evaluate the kinetics of T and the patterns of T recovery after short-term AD (STAD) and long-term AD (LTAD) when combined with high-dose external-beam radiotherapy (EBRT).

Methods. Between 2005 and 2010, 361 patients with intermediate and high-risk PCa were randomized to receive STAD (4 months) versus LTAD (28 months) combined with EBRT as part of a multicentre Spanish prospective randomized phase III trial (DART01/05). Study endpoints included overall survival, metastasis free survival, disease-free survival and biochemical disease-free survival. The serum T and PSA data were obtained every 3 months for the first 2 years and every six months thereafter. Castration level was defined as serum T < 0.5 ng/ml. Time to recovery of serum T level was measured from the last hormone treatment until the time when normal T values (> 1 ng/ml) were reached. The median follow-up was 42 months.

Results. Median T basal levels were 4.1 ng/ml and 4.0 ng/ml for patients in the LTAD and STAD arms, respectively. There was no significant correlation between T basal level and patient age, Gleason score, T stage, risk subgroup and biochemical failure (BF). The percentage of patients that experienced recovery of T normal values following ADT was 43.8% in LTAD group and 82.7% in the group treated with STAD. By the Kaplan–Meier method, the median time to recover T normal values was 9.8 months and 22 months ($p < 0.001$) for STAD and LTAD, respectively. The median time to BF was 60.2 months for LTAD (60.6 and 56.9 in patients with and without recovery of normal T values respectively). For STAD group, the median time to BF was 28 months (28.4 and 18.9 respectively).